

Journal of
Obstetric,
Gynecologic, &
Neonatal Nursing

45
1972 - 2016

May/June 2016

Volume 45 • Number 3

JOGNN

*Scholarship for the Care of Women,
Childbearing Families, & Newborns*



EDITORIAL

Progress Toward Meeting the Goals of
The Future of Nursing and Implications for
Maternity Care

REVIEW

An Integrative Review of Mothers'
Experiences of Preeclampsia

RESEARCH

The Lived Experience of Intimate Partner
Violence in the Rural Setting

Perceptions of Electronic Cigarettes
Among Medicaid-Eligible Pregnant and
Postpartum Women

Prepregnancy Obesity and a
Biobehavioral Predictive Model for
Postpartum Depression

Skin-to-Skin Care After Birth for
Moderately Preterm Infants

Predictors of Breastfeeding Initiation and
Frequency for Preterm Infants in the NICU

The Effects of Single-Family Rooms
on Parenting Behavior and Maternal
Psychological Factors

Barriers That Impede the Provision of Pain
Care to Neonates by Nurses in Jordan

Factors Associated With Perceived Health
Status of Multiracial/Ethnic Midlife Women
in the United States

SPECIAL REPORT

Current Resources for Evidence-Based
Practice, May/June 2016

IN FOCUS EDITORIAL

A Call to Action to Prevent
Cardiovascular Disease in Women

IN FOCUS CNE

Guidelines for the Reduction of
Cardiovascular Disease in Women

IN FOCUS

Pregnancy Reveals Evolving Risk for
Cardiometabolic Disease in Women

A Review of the Literature on Cardiac
Symptoms in Older and Younger Women

Subclinical Cardiovascular Disease in
Lymphoma Survivors by Sex

AWHONN POSITION STATEMENTS

Midwifery

Human Trafficking

Gender Bias in Women's Health,
Obstetric, and Neonatal Nursing



EDITORIAL

297 Progress Toward Meeting the Goals of *The Future of Nursing and Implications for Maternity Care*

The provision of quality maternity and women's health care for the burgeoning population of women in the United States depends on the ability of nurses to meet the IOM challenges.

Nancy K. Lowe

REVIEW

300 An Integrative Review of Mothers' Experiences of Preeclampsia

The experience of preeclampsia was characterized by fear, feeling close to death, and separation; these feelings changed to hope when the infant survived in the NICU.

Inger Emilie Værland, Kari Vevatne, and Berit Støre Brinchmann

RESEARCH

308 The Lived Experience of Intimate Partner Violence in the Rural Setting

Nurse management of intimate partner violence must be undertaken as part of a comprehensive public health approach.

Karen Roush and Ann Kurth

320 Perceptions of Electronic Cigarettes Among Medicaid-Eligible Pregnant and Postpartum Women

Pregnant women are initially attracted to electronic cigarettes as a harm reduction strategy, but they often return to traditional cigarettes in the postpartum period.

Amanda Fallin, Alana Miller, Sara Assef, and Kristin Ashford

326 Prepregnancy Obesity and a Biobehavioral Predictive Model for Postpartum Depression

The identification of biobehavioral pathways in the etiology of depression during the postpartum period may provide opportunities for preventive strategies that improve outcomes for mothers and infants.

Sharon L. Ruyak, Nancy K. Lowe, Elizabeth J. Corwin, Madalynn Neu, and Blake Boursaw

339 Skin-to-Skin Care After Birth for Moderately Preterm Infants

Early skin-to-skin contact in the delivery room for moderately preterm newborns may be feasible and safe.

Laila Kristoffersen, Ragnhild Stoen, Linda Frances Hansen, Janne Wilhelmsen, and Hakon Bergseng

346 Predictors of Breastfeeding Initiation and Frequency for Preterm Infants in the NICU

Gestational age, early physical contact at the time of birth, and mother's attitude are significant factors that predicted breastfeeding in the NICU.

Hannakaisa Niela-Vilén, Hanna-Leena Melender, Anna Axelin, Eliisa Löyttyniemi, and Sanna Salanterä

359 The Effects of Single-Family Rooms on Parenting Behavior and Maternal Psychological Factors

Mothers of infants in single-family room nurseries were present more, had higher rates of breastfeeding, and reported less stress than mothers of infants in open ward nurseries.

Rachelle Jones, Liz Jones, and Anne-Marie Feary

371 Barriers That Impede the Provision of Pain Care to Neonates by Nurses in Jordan

Improving neonatal pain care practices in Jordan requires an understanding of neonatal pain, pain assessment, and pain management options and effective interprofessional collaboration.

Nadin M. Abdel Razeq

378 Factors Associated With Perceived Health Status of Multiracial/Ethnic Midlife Women in the United States

The results indicated that researchers should consider differences among racial/ethnic groups when assessing women's perceived health status.

Young Ko, Wonshik Chee, and Eun-Ok Im

On Our Cover:

Vincent van Gogh. Sunflowers. 1889. Van Gogh Museum. Photo Credit: Album/Art Resource, NY.

Continued

SPECIAL REPORT

391 Current Resources for Evidence-Based Practice, May/June 2016

A review of new resources to support the provision of evidence-based care for women and infants.

Nicole S. Carlson

IN FOCUS EDITORIAL

400 A Call to Action to Prevent Cardiovascular Disease in Women

The guest editor suggests that precursors for cardiovascular disease that manifest during pregnancy present opportunities for further screening, monitoring, and lifestyle interventions for at-risk women.

Sandra B. Dunbar

IN FOCUS CNE

402 Guidelines for the Reduction of Cardiovascular Disease in Women

CNE Updated cardiovascular guidelines suggest important approaches for improving care of women to prevent cardiovascular disease, reduce risk, and ultimately improve cardiovascular outcomes in women.

Gina P. Lundberg, Sandra B. Dunbar, and Nanette K. Wenger

IN FOCUS

413 Pregnancy Reveals Evolving Risk for Cardiometabolic Disease in Women

Adverse pregnancy conditions such as hypertensive disorders of pregnancy, gestational diabetes, and preterm birth indicate elevated future risk for cardiovascular disease in women of childbearing age.

Erin P. Ferranti, Emily J. Jones, and Teri L. Hernandez

426 A Review of the Literature on Cardiac Symptoms in Older and Younger Women

Women older than 65 years reported less chest pain and more dyspnea and were more likely to experience prodromal symptoms before acute coronary syndrome.

Holli A. DeVon, Christina M. Pettey, Karen M. Vuckovic, Mary Dawn Koenig, and Jean C. McSweeney

438 Subclinical Cardiovascular Disease in Lymphoma Survivors by Sex

When compared with men in the 2 to 10 years after lymphoma treatment, women had greater than expected subclinical cardiovascular disease, lower health-related quality of life, diminished functioning, and more symptoms.

Carolyn Miller Reilly, Natia Esiashvili, Susmita Parashar, and Melinda Higgins

AWHONN POSITION STATEMENTS

454 Midwifery

458 Human Trafficking

461 Gender Bias in Women's Health, Obstetric, and Neonatal Nursing

ARTICLES ONLINE

Access these and all articles at <http://jognn.org>.

IN FOCUS CNE POST TEST

e5 Guidelines for the Reduction of Cardiovascular Disease in Women

Gina P. Lundberg, Sandra B. Dunbar, and Nanette K. Wenger